

OXALIDACEAE OXALIS FAMILY

Robert Ornduff
 Department of Integrative Biology
 University of California
 Berkeley, CA 94720
 and
 Melinda Denton
 Department of Botany
 University of Washington
 Seattle, WA 98195

Annual or perennial herbs, shrubs, or rarely trees. LEAVES alternate, palmately or pinnately compound (sometimes reduced to 1 leaflet), usually with petioles. INFLORESCENCES cymes, sometimes umbel- or raceme-like, or flowers solitary, usually axillary; peduncle bracted. FLOWERS generally perfect, actinomorphic, often heterostylous, with 5 distinct sepals and petals or these fused at base; stamens 10 or 15, connate below, usually of 2 lengths; pistil 1, with 3-5 carpels, the locules equalling carpels in number; styles 1-5, usually distinct. FRUITS usually fleshy loculicidal capsules. SEEDS usually arillate. --8 genera, 575 spp., chiefly of temperate regions.

Oxalis L. Oxalis, Wood-sorrel

In AZ perennial herbs with bulbs or creeping stems, caulescent or acaulescent. LEAVES petiolate, palmately compound, with 3 or more obcordate leaflets. FLOWERS perfect, the petals clawed, connate at base; stamens 10; pistil 5-carpelled; styles 5. FRUITS fleshy explosive capsules. SEEDS arillate. --ca. 500 spp., mostly of temperate regions and especially concentrated in s S. Amer. and s Afr. Many species are of considerable ornamental value, although several of these have become invasive weeds in many regions. (Greek: *oxus* = sour). Collections must include underground parts to facilitate identifications and labels should note fresh corolla color. Eiten, G. 1963. Amer. Midl. Naturalist 69:257-307; Lourteig, A. 1975. Phytologia 42:57-197; Denton, M. 1973. Pub. Museum Michigan State Univ., Biol. Series 4:459-615.

1. Plants without an above ground stem, with bulbs; corollas yellow, purple, blue, pink, or white.
2. Leaves arising mostly from underground stems; corollas yellow; [section *Cernuae*]
 *O. pes-caprae*
- 2' Leaves arising directly from bulbs; corollas purple, blue, pink, white, or intermediate shades (not yellow); [sect. *Ionoxalis*].
3. Flanges (conspicuous membranous margins) at petiole bases extended 0.5-1.5 cm above the bulb; leaflets 3-5, often with 2 or more oxalate deposits at the distal end of each lobe
 *O. caerulea*
- 3' Flanges at petiole bases barely, if at all, extended above the bulb; leaflets 3-11, with oxalate deposits at the notch, along the margins, randomly distributed, or absent.
4. Leaflets 5-11 on mature plants, usually 1.2 times or more longer than wide; bulb scales usually with 5 or more veins *O. decaphylla*
- 4' Leaflets 3 on mature plants, as wide or wider than long; bulb scales 3-veined.
5. Seeds with longitudinal ridges, rarely with faint transverse ridges, 0.8 mm wide or less; leaflet lobes to 1/5 leaflet length; bulblets frequently formed
 *O. alpina*
- 5' Seeds with longitudinal and transverse ridges, 0.8 mm wide or more; leaflet lobes to 1/4 or more leaflet length; bulblets absent *O. drummondii*

- 1' Plants with an above ground stem, without bulbs; corollas yellow or orange; [section *Corniculatae*]
6. Septate hairs present on stems, petioles, or pedicels; 1-3 erect stems arising from a slender underground rhizome *O. stricta*
- 6' Septate hairs absent from stems, petioles, or pedicels; few to many lax or prostrate stems arising from a thickened taproot.
7. Taproot woody; stem hairs often s-shaped *O. albicans*
- 7' Taproot fleshy; stem hairs, if present, straight or slightly curved.
8. Stem hairs absent, or, when present, straight or slightly curved; stipules rectangular; stems often rooting at nodes *O. corniculata*
- 8' Stem hairs straight; stipules oblong; stems not rooting at nodes *O. dillenii*

Oxalis albicans Kunth (tending to white). --BULBS absent. STEMS lax to 40 cm long, decumbent, at least at base, not rooting at the nodes. LEAVES cauline; leaflets 3, to 1.5 cm long, pale green. INFLORESCENCES 1-3-flowered; pedicel less than 2 cm long. COROLLAS yellow or orange, 8-12 mm long. FRUITS cylindric, 6-18 mm long. SEEDS 1.2-1.7 mm long. 3 subspp.; CA to TX, s to Guatemala.

Subsp. *albicans* --Stem and pedicel hairs crisped or straight, appressed or ascending; longer stem hairs 0.3-0.8 mm long. Desert scrub, oak, pine, or oak-pine woodland: Gila, Graham, (La Paz?), Maricopa, Mohave, Pima, Pinal, Santa Cruz, Yavapai, Yuma cos.; 750-2000 m (2200-6000 ft); Feb-Oct; w TX to Baja C. Sur, Mex.; Guatemala.

Subsp. *pilosa* (Nuttall) Eiten (pilose). --Stem and pedicel hairs straight, appressed, spreading, or retrorse; longer stem hairs 0.7-1.2 mm long. [*O. pilosa* Nuttall]. --Chaparral, desert woodland, oak or mixed deciduous woodland, or occasionally weedy in urban areas: Cochise, Coconino, Gila, Graham, (La Paz?), Maricopa, Mohave, Pima, Pinal, Santa Cruz, Yavapai, Yuma cos.; 600-2200 m (1900-6600 ft); Feb-Oct; CA to TX; n Mex.

Oxalis alpina (Rose) Knuth (alpine). --BULBS 0.8-2 cm long, often with up to 20 bulblets. LEAVES 4-30 cm tall; leaflets 3, 4-27 mm long, 4-38 mm wide. INFLORESCENCES 1-7-flowered; pedicels less than 5 cm long. COROLLAS blue, lavender, pink, or white, to 27 mm long. FRUITS ellipsoid, 5-12 mm long. SEEDS 1-1.5 mm long. [*O. metcalfei* Small]. $2n = 14, 28, 42, 56, 84$. --Open, moist, often rocky places in deciduous, pine-oak, or coniferous forests: Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Pima, Pinal, Santa Cruz, Yavapai cos.; 1700-3200 m (5000-9600 ft); Jul-Sep; NM, TX, s to Guatemala.

Oxalis caerulea (Small) Knuth (dark blue). --BULBS 0.8-1.5 cm long. LEAVES 3-14 cm tall; leaflets 3-5, 5-23 mm long, 5-22 mm wide. INFLORESCENCES 1-7-flowered; pedicels less than 15 mm long. COROLLAS rose-red or lavender, 3.5-14 mm long. FRUITS ellipsoidal, 3.5-6 mm long. SEEDS 1 mm long. --Pine forests: Apache, Coconino, Yavapai cos.; 2300-3000 m (6900-9000 ft); Jul-Sep; NM; s to Dgo., Mex.

Oxalis corniculata L. (with small curved horn-like appendages). --BULBS absent. STEMS prostrate or decumbent, to 30 cm long, often rooting at nodes. LEAVES cauline; leaflets 3, to 2 cm long, often maroon. INFLORESCENCES 2-5-flowered; pedicels less than 1 cm long. COROLLAS yellow, 4-8 mm long. FRUITS cylindric, angled, 6-25 mm long. SEEDS 1.2-1.5 mm long. $2n = 24, 28, 44, 48$. -- Common in lawns, gardens, disturbed places, greenhouses, and occasionally woodlands or grasslands: Apache, Gila, Maricopa, Mohave, Pima, Santa Cruz cos.; 250-1300 m (1100-3900 ft); flowers all year; introduced from the Old World.

Oxalis decaphylla HBK (ten leaflets). Agritos. --BULBS 1-3 cm long. LEAVES 6-24 cm tall; leaflets 5-11, 10-72 mm long, 2-40 mm wide. INFLORESCENCES 5-12-flowered; pedicels less than 30 mm long. COROLLAS purplish, pink, or lavender, 7-22 mm long. FRUITS ellipsoid, 3-11 mm long. SEEDS 0.8-1.2 mm long. [*O. grayi* Rose]. $2n = 28, 56$. --Coniferous, oak, or mixed deciduous forests, sometimes

scrubland or grassland; occasionally weedy: Apache, Cochise, Coconino, Gila, Greenlee, Navajo, Pima, Santa Cruz, Yavapai cos.; 1700-3200 m (5000-9500 ft); Jul-Sep; NM to Mex.

Oxalis dillenii Jacquin (for J. J. Dillenius). --BULBS absent. STEMS erect or decumbent, to 40 cm long. LEAVES cauline; leaflets 3, 4-18 mm long, pale green. INFLORESCENCES 1-5-flowered; pedicels 10-30 mm long. COROLLAS yellow, 5-12 mm long. FRUITS cylindrical or prismatic, 8-25 mm long. SEEDS 1-1.5 mm long. $2n = 16, 18, 20, 22$. --Moist areas, grassland, coniferous forests: Apache, Coconino, Gila, Graham cos.; 2100-2500 m (6300-7500 ft); May-Sep; s Can; e US to WA, MT to NM, and TX.

Whether the AZ plants are correctly assigned to this species or are more correctly referred to *O. priceae* Small is unclear.

Oxalis drummondii Gray (for T. Drummond). --BULBS 1-2 cm long. LEAVES 6-20 cm tall; leaflets 3, 6-34 mm long, 10-50 mm wide. INFLORESCENCES 3-10-flowered; pedicels less than 35 mm long. COROLLAS pink or lavender, 8-20 mm long. FRUITS cylindrical, 4-12 mm long. SEEDS 1-1.5 mm long. [*O. amplifolia* (Trelease) Knuth in part]. $2n = 14$. Scrublands or coniferous forests: Cochise, Santa Cruz cos.; 1200-2700 m (3500-8000 ft); Aug; TX, s to Tamp., Mex. Many AZ specimens variously exhibit features of *O. alpina*, *O. latifolia*, and *O. violacea* and require further study.

Oxalis pes-caprae L. (foot of a goat). Bermuda Buttercup. --BULBS 0.5-2.5 cm long. LEAVES 3-42 cm tall; leaflets 3, 10-40 mm long, 20-45 mm wide. INFLORESCENCES 3-20 flowered; pedicels less than 2 cm long. COROLLAS yellow, 15-25 mm long. FRUITS and SEEDS unknown in N. Amer. specimens. $2n = 14, 28, 35$. --Disturbed urban or agricultural lands: Maricopa, Pima, Yavapai cos.; 400-650 m (1100-2000 ft); Feb- Apr; introduced from South Africa.

In its native South Africa this species is represented by tristylous populations, but in N. Amer. only a pentaploid, sterile, short-styled morph is known.

Oxalis stricta L. (very upright). -- BULBS absent. STEMS usually erect, unbranched, to 75 cm tall. LEAVES cauline; leaflets 3, 5-20 mm long, pale green. INFLORESCENCES 1-9, rarely more, 2-5-flowered; pedicels 8-10 cm long. COROLLAS yellow, or orange-yellow, 3.5-11 mm long. FRUITS cylindrical, 8-13 mm long. SEEDS 0.8-1.3 mm long. $2n = 18, 24$. Woodlands: Coconino, Maricopa cos.; 2100-2200 m (6300-6700 ft); Apr-Jul; s Can. to se U.S., scattered in c and sw U.S.; e Asia; introduced elsewhere.

This treatment was prepared by Ornduff, and was adapted from Eiten's revision of sect. *Corniculatae*, from Denton's 1973 monograph of sect. *Ionoxalis* and her unpublished 1974 treatment of that group for the projected Chihuahuan Desert Flora, and from specimens at ARIZ, ASC, ASU, and MNA. Denton reviewed and slightly revised the sect. *Ionoxalis* treatment that appears here, before her death in 1994. In respect to AZ plants, the treatments of sect. *Corniculatae* published by Eiten and by Lourteig differ from each other in several important respects. The disagreements cannot be resolved without a worldwide study of the section using modern methods. Our treatment must be regarded as a tentative and provisional one. Because specific differences in sect. *Corniculatae* are unclear, only a map for the section has been made.




Figure 1. A, *Oxalis decaphylla* (reproduced from M. Denton's thesis). B, *Oxalis albicans* ssp. *pilosa* (reproduced with permission from The Jepson Manual, University of California Press).




Figure 2. Distribution of: A, *Oxalis alpina* (dots), *O. caerulea* (triangles); B, *O. decaphylla* (dots), *O. drummondii* (triangles); C, *Oxalis* section *Corniculatae* (including *O. albicans*, *O. corniculata*, *O. dillenii*, and *O. stricta*).